

## Review Of Research

### Abstract:-

This paper present the air pollution emitted from an Industrial area Satna. Air quality of SatnaIndustrial area was monitored for two successive years i.e. 2011-2012 and 2012-2013 at various sites for RSPM, SO<sub>x</sub> NO<sub>x</sub> &CO .The data collected were compared with Indian Standard prescribed by CPCB. Sampling of air quality was done at various distances from the industrial area at six sampling points. It was observed that SPM was higher than the permissible limit at all the sampling points except. Air quality index was also calculated and this study indicates that the air quality of Satna is unhealthy at various sampling site. People with asthma or other respiratory diseases, the elderly, and children are the groups most at risk.

### Keywords:

Air pollution, Industrial area Satna, PM<sub>10</sub>, RPM, SO<sub>2</sub>, NO<sub>x</sub> and CO.

**Shailendra Singh<sup>1</sup> , M. K. Bhatnagar<sup>2</sup> , Rajesh Sharma<sup>3</sup> and D.P. Prajapati<sup>4</sup>**

<sup>1</sup>Research Scholar Department of Chemistry Pt. S.N.S. Govt. P.G. College Shahdol (M.P.)

<sup>2</sup>Prof. & Head Department of Chemistry Pt. S.N.S. Govt. P.G. College Shahdol (M.P.)

<sup>3</sup>Department of Chemistry Govt. Chhatrashal P.G. College, Panna (M.P.)

<sup>4</sup>Asst. Prof. Department of Chemistry Govt. P.G. College Seoni (M.P.)

## STUDIES ON AIR QUALITY OF SATNA INDUSTRIAL AREA(M.P.)



## INTRODUCTION

Environment means all the surrounding conditions, which influence the growth and development of life. The environment is composed of gases and substances, which exist in a certain proportion. When for any reason, one of the constitutions increases or decreases the balance of the whole atmosphere is disturbed. This imbalance of the constituents is called pollution of the environment. Pollution is the introduction of contaminants into an environment that causes instability, disorder, harm or discomfort to the ecosystem i.e. physical systems or living organisms. Pollution can take the form of chemical substances, or energy, such as noise, heat or light energy. Pollutants, the elements of pollution, can be foreign substances or energies, or naturally occurring; when naturally occurring, they are considered contaminants when they exceed natural levels. Pollution is often classed as point source or non-point source pollution. The pollution is chiefly two types - Air and water pollutions.

Today we breathe in the industrials pollution every moment. On the other hand we add that in one way and the other. We have slums cropping up like mescal in tops side by side the sooty chimneys and take pride in our industrial progress.

Pollution means of the addition of any foreign material (Inorganic, biological, or radiological) or any physical change in the natural water, which may harmfully affecting the living life (human, agricultural or biological) directly or indirectly. After some times or very long times.

According to general thinking of people "pollution means the introduction in to natural water of anything that to them appears to be foreign. The addition of something to water which changes its natural quality.

The environmental pollution is the unfavorable alternation of our surrounding. Wholly are largely as a byproduct of mains action. Through direct or indirect effect of changes in energy patterns. Radiation levels chemical and physical constitution and abundance of organisms.

Although water pollution is an age old problem but in this modern age. The problems like population increase sewage disposal industrials waste radioactive waste are some heavy metals etc. We have polluted our water resources so much so that about 70% rivers and streams not only of India but of all the countries contain polluted waters. We will discuss some important sources of industrial pollution along with the constituents which pollute the rivers stream and fields.

The rapid industrialization leading to urbanization, unplanned and excessive exploitation of natural resources have been causing pollution problems in cities and towns of developing countries. Manmade and natural sources of emissions have polluted the air with toxic substances. The national average per capita SO<sub>x</sub> emission was 4.2 kg per person in 1990, which rose to 5 kg in 1995, an increase of almost 20% in 5 yr. In 1990, coal consumption contributed 64% of total SO<sub>x</sub> emissions in India, oil products 29%, biomass 4.5% and non-energy consumption 2.5% (Garget, al, 2001). Total SO<sub>x</sub> and NO<sub>x</sub> emission from India were 3542 and 2636 respectively (1990) and 4638 and 3462 (1995) growing at annual rate of around 5.5%. The sectoral composition of SO<sub>x</sub> emissions indicates a predominance of electric power generation sector [46%]. Power and transport sector emissions equally dominate NO<sub>x</sub> emissions contributing nearly 30% each (Garget, al, 2001). A recent study in middle-income homes of Delhi found PM<sub>10</sub> levels to be as high as 170–810 µg/m<sup>3</sup> even in homes where there was no cooking or smoking activity (Kumar, 2001).

A report showed that SPM concentrations in Shanghai, New Delhi, Mumbai, Guangzhou, Chongquin, Calcutta, Beijing and Bangkok exceeded WHO limits (90 µg/m<sup>3</sup>) by three, five, three, three, four, four, four and two times respectively. It also showed that PM<sub>10</sub> exceeded the USEPA limit (50 µg/m<sup>3</sup>) by several times in a number of cities, most notably by over four times in New Delhi and Calcutta. Data from Tokyo shows that TSP increased rapidly from 40 µg/m<sup>3</sup> in the early 1980 to over 70 µg/m<sup>3</sup> in the early 1990; after that TSP has been decreasing or stagnating, but it is becoming an increasing challenge to contain TSP and NO<sub>x</sub>. (Saksena et al, 2003).

A study for assessment and management of air quality was carried out in the Ib Valley area of the Ib Valley coalfield in Orissa state, India. The 24 h average concentrations of total suspended particulate (TSP), respirable particulate matter (PM<sub>10</sub>), sulfur dioxide (SO<sub>2</sub>) and oxides of Nitrogen (NO<sub>x</sub>) were determined at regular intervals throughout one year at twelve monitoring stations in residential areas and six monitoring stations in mining/industrial areas. The 24 h average TSP and PM<sub>10</sub> concentrations were 124.6–390.3 µg/m<sup>3</sup> and 25.9–119.9 µg/m<sup>3</sup> in residential areas, and were 146.3–845.2 µg/m<sup>3</sup> and 45.5–290.5 µg/m<sup>3</sup> in industrial areas. During the study period, 24 h and annual average TSP and PM<sub>10</sub> concentrations exceeded the respective standards set in the Indian national ambient air quality standard (NAAQS) protocol as well as USEPA, EU, WHO and World Bank standards at most of the residential and industrial areas. However, concentrations of SO<sub>2</sub> (annual average: 24.6–36.1 µg/m<sup>3</sup> and 24 h, average: 17.0–46.3 µg/m<sup>3</sup>) and NO<sub>x</sub> (annual average: 23.6–40.9 µg/m<sup>3</sup> and 24 h average: 18.3–53.6 µg/m<sup>3</sup>) were well within the prescribed limit of the NAAQS and international standards in both residential and industrial areas (Chaulaya, 2004).

Monitoring of ambient respirable suspended particulate matter or PM<sub>10</sub> and total suspended particulate (TSP) levels around a large coal-fired power station in India was carried out. Geometric mean values of PM<sub>10</sub> and TSP concentrations at sites of downwind direction are found in ranges of 74–144 µg/m<sup>3</sup> (PM<sub>10</sub>) and 459–647 µg/m<sup>3</sup> (TSP) in post rainy season, 154–207 µg/m<sup>3</sup> (RSPM) and 437–610 µg/m<sup>3</sup>

(SPM) in summer season and 180–275  $\mu\text{g}/\text{m}^3$  (PM10) and 933–1578  $\mu\text{g}/\text{m}^3$  (TSP) in winter season (Sharma et. al, 2005).


The rural-industrial site at Satna shows significant different from urban, urban-industrial, rural, rural-remote and rural–urban influenced sites. With a minimum of one daily sample at each site, mean PM10 concentration at the rural-industrial Satna site varied from 65.5 to 147.5  $\mu\text{g}/\text{m}^3$ , and from 205.0 to 320.3  $\mu\text{g}/\text{m}^3$  at the urban-industrial Delhi site. With a minimum of one daily sample at each site, the mean TSP concentration at the rural-industrial Satna site varies from 283.9 to 678.0  $\mu\text{g}/\text{m}^3$ , while at the urban industrial Delhi site mean TSP concentration varies from 553.4 to 827.6  $\mu\text{g}/\text{m}^3$ . The maximum TSP concentration in Satna, 678.0  $\mu\text{g}/\text{m}^3$ , was on December 9, 2000. Maximum TSP concentration in Delhi, 827.6  $\mu\text{g}/\text{m}^3$ , occurred on October 23, 2000. Levels of TSP in Ashok Vihar, a residential area in Delhi, reached 10 times the permissible limit in October 2000 (Kaushiket, al, 2007).

### STUDY AREA-

Satna is located at 24.34°N 80.55°E with an elevation of 315 meters in central India and a municipal corporation in Satna district in the Indian state of Madhya Pradesh. Satna is a border city of the state and is touched by the border of state of Uttar Pradesh. Satna district has Uttar Pradesh on the north. Rewa on the east Shahdol on the southwest. Umaria and Katni on the south & Panna on the west. The District is part of Rewa Division, the Satna District is divided into the Tehsil of Amarpatan, Maihar, Nagod and Raguraj Nagar. Satna town is situated in Raghuraj Nagar Tehsil, the place is renowned for dolomite mines & limestone.

Satna is in the limestone belt of India. It contributes around 8-9 % of India Total cement production. There is abundance of dolomite and limestone in the area, the city has seven cement factories that are producing cement and exporting the same other part of the country. The electrical cable factory “Universal Cables” in Satna is among the pioneers in the country. The city of Satna is known as commercial capital of Baghelkhand. The city is amongst few most promising cities of Madhya Pradesh because of several new industries as- Iron industries, plywood factory, Detergent factory, Stone crusher, medical hospital, transport auto workshop electrical motor & submersible manufacturing diesel rail engine workshop heavy motor workshop & other new planned by some of the reputed industrial houses in the country. Present major problem faced in city may inter alia include electricity, water pollution, air pollution due to atmospheric wastes of cement factories. Satna is known as cement city in India. Due to abundance of limestone and dolomite in the region.

The study area is surrounded by Cement factories. Due to these industries the study area is under immense Environmental stress.



### MATERIAL AND METHOD:

The present work is aimed to assess the Environmental status of the Satna industrial area. The study is focused only on two parameters – Air quality monitoring parameters. Based on both parameters,

well versed methodology is made separately for each parameter. Standard methods of air analysis prescribed by CPCB standard methods prescribed by APHA has been used.

### RESULTS & DISCUSSION :

In order access air quality of Satna Industrial area ambient air monitoring have been done. For this purpose air pollutant RSPM, SO<sub>x</sub>, NO<sub>x</sub> & CO in ambient air of study area has been analysed.

To assess the air pollution in Satna Industrial area various sampling site for ambient air analysis were selected. The different sampling stations are given below –

- 1.Santoshi Mata Mandir (A1): App. 3 Km. Distance from Semariya Chowk.
- 2.Sagmaniha (A2): App. 3.5 Km. Distance from Semariya Chowk.
- 3.Gaharwar petrol Pump (A3): App. 4 Km. Distance from Semariya Chowk.
- 4.Biruhali Railway Station (A4): App. 6 Km. Distance from Semariya Chowk.
- 5.Raj Stone Crusher (A5): App. 7 Km. Distance from Semariya Chowk.
- 6.Semariya Gate (A6): App. 8 Km. Distance from Semariya Chowk.

Air quality of Satna Industrial area was monitored for two successive years i.e. 2011-2012 and 2012-2013 at various sites for RSPM, SO<sub>x</sub>, NO<sub>x</sub> & CO.

The various results are given in Tables. These parameters are regularly monitored in each season in every year.

At Santoshi Mandir site RSPM was found to be above the Permissible limit in almost every season in the year 2011 and 2012 except in monsoon in 2011-12 when its value was recorded 115 µg/m<sup>3</sup>. Its range was from 115 µg/m<sup>3</sup> to 140 µg/m<sup>3</sup> (2011-12) and 135 µg/m<sup>3</sup> to 143 µg/m<sup>3</sup> (2012-13), Sulphur dioxide and oxides of nitrogen were found to be within permissible limit. The concentration of SO<sub>x</sub> was found to be in between 13 µg/m<sup>3</sup> to 20.4 µg/m<sup>3</sup> (2011-12) & 16 µg/m<sup>3</sup> to 16.7 µg/m<sup>3</sup> (2012-13), Similarly No<sub>x</sub> was observed in the range between 21.6 µg/m<sup>3</sup> to 24.6 µg/m<sup>3</sup> (2011-12) & 21.5 µg/m<sup>3</sup> to 24 µg/m<sup>3</sup> (2012-13). The CO concentration falls within the ranges of 1 to 1.8 µg/m<sup>3</sup> (2011-12) & 1 to 1.7 µg/m<sup>3</sup> in 2012-13.

In 2011-12 Air quality at Sagmaniha which is situated about 3.5 Km. from Semariya Chowk was found to be within the Permissible limit except RSPM value. At this site RSPM was assessed in the range of 129 µg/m<sup>3</sup> to 132 µg/m<sup>3</sup> and Value of SO<sub>x</sub> was observed in between 14.2 µg/m<sup>3</sup> to 20.6 µg/m<sup>3</sup>, while NO<sub>x</sub> was range between 23.3 µg/m<sup>3</sup> to 27 µg/m<sup>3</sup>. The concentration of CO was recorded in the range of 1.1 to 1.9 µg/m<sup>3</sup>.

While in 2012-13 at this site RSPM was assessed in the range of 132 µg/m<sup>3</sup> to 140 µg/m<sup>3</sup> and Value of SO<sub>x</sub> was recorded from 15.8 µg/m<sup>3</sup> to 25 µg/m<sup>3</sup>. No<sub>x</sub> was observed in between 21 µg/m<sup>3</sup> to 24.5 µg/m<sup>3</sup> while CO was in the range between 1 µg/m<sup>3</sup> to 1.8 µg/m<sup>3</sup>.

During the pollution assessment of Sagmaniha monitoring station. It indicates the higher trend of Respirable suspended Particulate matter dispersion into surrounding environment especially in 2012 in respect to 2011 year because of heavy transportation in nearby area.

**Table – 1 Different site in air sample year 2011-12**

Sample Site	Unit	So <sub>x</sub>			No <sub>x</sub>			CO			RSPM		
		W	S	M	W	S	M	W	S	M	W	S	M
A-I	µg/m <sup>3</sup>	15	13	20.4	21.6	23.9	24.6	1.8	1	1.6	140	120	115
A-II		14.2	14.0	20.6	23.3	24.7	27	1.10	1.4	1.9	132	130	129
A-III		15.5	13.5	20.2	20.1	23	23.9	1.9	1.4	1.5	142	134	121
A-IV		14.5	13.5	20.1	24.6	24.7	20.9	1.1	1.10	2	140	125	120
A-V		16.5	15	20	25.9	24.3	25.6	1.10	1.8	1.8	119	121	120
A-VI		16.0	14.5	19.5	20.1	23	26.6	1.15	1.6	1.10	149	135	125

W= Winter S= Summer = Monsoon

**Table – 2 Different site in air sample year 2012-13**

Sample Site	Unit	So <sub>x</sub>			No <sub>x</sub>			CO			RSPM		
		W	S	M	W	S	M	W	S	M	W	S	M
A-I	µg/m <sup>3</sup>	15	13	20.4	21.6	23.9	24.6	1.8	1	1.6	140	120	115
A-II		14.2	14.0	20.6	23.3	24.7	27	1.10	1.4	1.9	132	130	129
A-III		15.5	13.5	20.2	20.1	23	23.9	1.9	1.4	1.5	142	134	121
A-IV		14.5	13.5	20.1	24.6	24.7	20.9	1.1	1.10	2	140	125	120
A-V		16.5	15	20	25.9	24.3	25.6	1.10	1.8	1.8	119	121	120
A-VI		16.0	14.5	19.5	20.1	23	26.6	1.15	1.6	1.10	149	135	125

Sample Site	So <sub>x</sub>			No <sub>x</sub>			CO			RSPM		
	W	S	M	W	S	M	W	S	M	W	S	M
A-I	16.7	16.2	16	24.2	21.5	21.7	1.10	1.7	1	143	141	135
A-II	25	15.8	15.9	24.5	22.7	21	1.8	1	1.7	140	138	132
A-III	18	17.7	17	24.4	21	21.10	1.7	1.4	1	139	136	132
A-IV	17.5	17	16.9	29.7	19	19.9	1.2	0	1.2	141	140	136
A-V	17	16.3	16.9	20.7	20	20.1	1	0	1	136	130	128
A-VI	20.9	20	18.7	23	22.9	20.5	1.20	1.15	1.10	142	140	140

In 2011 at Gaharwar petrol pump which is situated at 4 Km distance from Semariyachowk, the RSPM was found to be higher than the permissible limit. The concentration range of RSPM was from 121  $\mu\text{g}/\text{m}^3$  to 142  $\mu\text{g}/\text{m}^3$  and at this site Sox was assessed in the range of 94.36  $\mu\text{g}/\text{m}^3$  to 156.42  $\mu\text{g}/\text{m}^3$ . The concentration of SO<sub>x</sub> and NO<sub>x</sub> was found in the range of 13.5  $\mu\text{g}/\text{m}^3$  to 20.2  $\mu\text{g}/\text{m}^3$  and 20.1  $\mu\text{g}/\text{m}^3$  to 23.9  $\mu\text{g}/\text{m}^3$ .

While in 2012 the RSPM was found to be higher than permissible limit at bamboo gate. The concentration range of R.S.P.M was from 135  $\mu\text{g}/\text{m}^3$  to 143  $\mu\text{g}/\text{m}^3$  and at this site Sox was assessed in the range of 16  $\mu\text{g}/\text{m}^3$  to 16.7  $\mu\text{g}/\text{m}^3$ . The concentration of No<sub>x</sub> & CO was found in the range of 21.7  $\mu\text{g}/\text{m}^3$  to 24.2  $\mu\text{g}/\text{m}^3$  and 1  $\mu\text{g}/\text{m}^3$  to 1.7  $\mu\text{g}/\text{m}^3$ .

The RSPM value at winter season in 2011-12 & 2012-13 was recorded maximum i.e. 142 & 143 respectively. At the same season both values are higher than the permissible limit. It can be due to weather condition and Vehicular activities at the site.

In 2011 The concentration of RSPM at Biruhali Railway Station (6 Km.) was found to be in the range 120  $\mu\text{g}/\text{m}^3$  to 140  $\mu\text{g}/\text{m}^3$ , which is above the permissible limit as prescribed by CPCB. The range of Sox was found to be in between 13.5  $\mu\text{g}/\text{m}^3$  to 20.1  $\mu\text{g}/\text{m}^3$ , while the range of No<sub>x</sub> was in between 20.9  $\mu\text{g}/\text{m}^3$  to 24.7  $\mu\text{g}/\text{m}^3$ . The value of CO was 1.1 to 2  $\mu\text{g}/\text{m}^3$ . So values of all these parameters fall within the permissible limit as prescribed by Central Pollution Control Board & M.P.P.C.B.

While in 2012 The value of R.S.P.M was in the range of 136  $\mu\text{g}/\text{m}^3$  to 141  $\mu\text{g}/\text{m}^3$ . The range of Sox was found to be in between 16.9  $\mu\text{g}/\text{m}^3$  to 17.5  $\mu\text{g}/\text{m}^3$ , While the range of No<sub>x</sub> was in between 19  $\mu\text{g}/\text{m}^3$  to 29.7  $\mu\text{g}/\text{m}^3$  while the value of CO ranges from 0 to 1.2  $\mu\text{g}/\text{m}^3$ . So values of all these parameters except RSPM fall within the permissible limit as prescribed by Central Pollution Control Board & M.P.P.C.B.

Dispersion of air pollutant is depend on wind speed, wind direction and location of monitoring stations, The monitoring station selected for the present study, is also located at North direction. which is most of the time, wind direction facing point therefore in 2012-13 results are comparatively higher than 2011-12 study i.e. RSPM values due to unlocalized distribution of air born dust, whereas Sox, No<sub>x</sub> parameter are not effected on the monitoring studies.

At Raj stone crushersite RSPM was found to be above the Permissible limit in almost every season in the year 2011 and 2012 except in winter in 2011-12 when its value was recorded 119  $\mu\text{g}/\text{m}^3$ . Its range was from 119  $\mu\text{g}/\text{m}^3$  to 121  $\mu\text{g}/\text{m}^3$  (2011-12) and 128  $\mu\text{g}/\text{m}^3$  to 136  $\mu\text{g}/\text{m}^3$  (2012-13), Sulphur dioxide and oxides of nitrogen were found to be within permissible limit. The concentration of SO<sub>x</sub> was found to be in between 15  $\mu\text{g}/\text{m}^3$  to 20  $\mu\text{g}/\text{m}^3$  (2011-12) & 16.3  $\mu\text{g}/\text{m}^3$  to 17  $\mu\text{g}/\text{m}^3$  (2012-13), Similarly No<sub>x</sub> was observed in the range between 20.9  $\mu\text{g}/\text{m}^3$  to 24.7  $\mu\text{g}/\text{m}^3$  (2011-12) & 20  $\mu\text{g}/\text{m}^3$  to 20.7  $\mu\text{g}/\text{m}^3$  (2012-13). The value of CO was 1.1 to 1.8  $\mu\text{g}/\text{m}^3$  (2011-12) & 0 to 1  $\mu\text{g}/\text{m}^3$  in 2012-13.

In 2011 The concentration of RSPM at Semariya Gate (8 Km. from semariyachowk) was found to be in the range 125  $\mu\text{g}/\text{m}^3$  to 149  $\mu\text{g}/\text{m}^3$ , which is above the permissible limit as prescribed by CPCB. The range of Sox was found to be in between 16.0  $\mu\text{g}/\text{m}^3$  to 19.5  $\mu\text{g}/\text{m}^3$ , while the range of No<sub>x</sub> was in between 20.1  $\mu\text{g}/\text{m}^3$  to 26.6  $\mu\text{g}/\text{m}^3$ . The value of CO was 1.1 to 1.6  $\mu\text{g}/\text{m}^3$ . So values of all these parameters fall within the permissible limit as prescribed by Central Pollution Control Board & M.P.P.C.B.

While in 2012 the value of R.S.P.M was in the range of 140  $\mu\text{g}/\text{m}^3$  to 142  $\mu\text{g}/\text{m}^3$ . The range of Sox was found to be in between 18.7  $\mu\text{g}/\text{m}^3$  to 20.9  $\mu\text{g}/\text{m}^3$ , While the range of No<sub>x</sub> was in between 20.5  $\mu\text{g}/\text{m}^3$  to 23  $\mu\text{g}/\text{m}^3$ . The value of CO ranges from 1.1 to 1.2  $\mu\text{g}/\text{m}^3$ . So values of all these parameters except RSPM fall within the permissible limit as prescribed by Central Pollution Control Board & M.P.P.C.B.

#### ACKNOWLEDGEMENT-

The authors are thankful to Dr. Sandeep Kumar Shukla, Asst. Prof., Department of zoology, Govt. P.G. College, Seoni (M.P.)



## REFERENCES

- 1.APHA, Standard Method for Examination of Water and Wastewater, 18th ed., American Public Health Association, Washington DC (1992).
- 2.Chaulya, S.K., Spatial and temporal variations of SPM, RPM, SO<sub>2</sub> and NO<sub>x</sub> concentrations in an opencast coal mining area, J. Environ. Monit., 6: 134-142(2004).
- 3.Garg A., Shukla, P.R., Bhattacharya, S. and Dadhwal, V.K., Sub-region (district) and sector level SO<sub>2</sub> and NO<sub>x</sub> emissions for India: Assessment of inventories and mitigation flexibility, Atmospheric Environment, 35, 703-713 (2001).
- 4.Kausik, S., Khare, M. and Gupta, K. B., Suspended particulate matter distribution in rural-industrial Satna and in urban-industrial south Delhi, Environ Monit. Assess, 128: 431–445(2007).
- 5.Kumar, A., Air Pollution Control in the Transportation Sector. Third Phase Research Report of the Urban Environmental Management Project, 2001.
- 6.Saksena, S., Singh, P.B., Prasad, R. K., Prasad, R., Malhotra, P., Joshi, V. and Patil, R.S., Exposure of infants to outdoor and indoor air pollution in low-income Urban areas - a case study of Delhi, Journal of Exposure Analysis and Environmental Epidemiology, 13,: 219-230 (2003).
- 7.Sharma, R., Pervez, Y. and Pervez, S., Seasonal variation and spatial variability of suspended particulate matter in the vicinity of a large coal-fired powerstation in India-A case study, Environmental Monitoring and Assessment, 102: 1-13 (2005).
- 8.Thareja S., S. choudhary and P. Trivedi Advances in Applied Research 2 (5),84. 2011.